

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
6 January 2005 (06.01.2005)

PCT

(10) International Publication Number  
**WO 2005/000835 A1**

- (51) International Patent Classification<sup>7</sup>: **C07D 401/12** (74) Agent: GRUBB, Philip; Novartis AG, Corporate Intellectual Property, CH-4002 Basel (CH).
- (21) International Application Number: PCT/EP2004/006915 (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (22) International Filing Date: 25 June 2004 (25.06.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/482,686 26 June 2003 (26.06.2003) US
- (71) Applicant (for all designated States except AT, US): NO-VARTIS AG [CH/CH]; Lichtstrasse 35, CH-4056 Basel (CH).
- (71) Applicant (for AT only): NOVARTIS PHARMA GMBH [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): SLADE, Joel [US/US]; 14 Theresa Drive, Flanders, NJ 07836 (US). VIVELO, James, Anthony [US/US]; 27 Vanderveer Drive, Basking Ridge, NJ 07920 (US). CHEN, Guang-Pei [US/US]; 33 Preston Drive, Livingston, NJ 07039 (US). BAJWA, Joginder, Singh [US/US]; 180D Donor Avenue, Elmwood Park, NJ 07407 (US). PARKER, David, John [US/US]; 291 E. Northfield Road, Livingston, NJ 07039 (US).
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PROCESS FOR PREPARING INTERMEDIATES USEFUL TO PREPARE CERTAIN ANTIBACTERIAL N-FORMYL HYDROXYLAMINES

(57) Abstract: The present invention is directed to a process for preparing intermediates that are useful to prepare certain antibacterial N-formyl hydroxylamine compounds which are peptide deformylase inhibitors.

WO 2005/000835 A1

BEST AVAILABLE COPY

**PROCESS FOR PREPARING INTERMEDIATES USEFUL TO PREPARE CERTAIN  
ANTIBACTERIAL N-FORMYL HYDROXYLAMINES.**


This invention is directed to a process for preparing intermediates that are useful to prepare certain antibacterial *N*-formyl hydroxylamine compounds.

Peptide deformylase is a metallopeptidase found in prokaryotic organisms such as bacteria. Protein synthesis in prokaryotic organisms begins with *N*-formyl methionine (fMet). After initiation of protein synthesis, the formyl group is removed by the enzyme peptide deformylase (PDF); this activity is essential for maturation of proteins. It has been shown that PDF is required for bacterial growth. See Chang et al., *J. Bacteriol.*, Vol. 171, pp. 4071-4072 (1989); Meinel et al., *J. Bacteriol.*, Vol. 176, No. 23, pp. 7387-7390 (1994); Mazel et al., *EMBO J.*, Vol. 13, No. 4, pp. 914-923 (1994). Since protein synthesis in eukaryotic organisms does not depend on fMet for initiation, agents that will inhibit PDF are attractive candidates for development of new anti-microbial and anti-bacterial drugs.

Co-pending Application Serial No. 10/171,706, filed June 14, 2002 (incorporated herein by reference in its entirety), PCT equivalent published as WO 02/102790 A1, discloses novel *N*-formyl hydroxylamine compounds that inhibit PDF and are therefore useful as antibacterial agents. The compounds disclosed therein are certain *N*-[1-oxo-2-alkyl-3-(*N*-hydroxyformamido)-propyl]-(carbonylamino-aryl or -heteroaryl)-azacyclo<sub>4,7</sub>alkanes or thiazacyclo<sub>4,7</sub>alkanes which are described in more detail hereinafter. An improved process has been discovered for preparing intermediates useful for preparing these *N*-[1-oxo-2-alkyl-3-(*N*-hydroxyformamido)-propyl]-(carbonylamino-aryl or -heteroaryl)-azacyclo<sub>4,7</sub>alkanes or thiazacyclo<sub>4,7</sub>alkanes.


The present invention is directed to a novel process for preparing certain intermediates which are useful to prepare certain *N*-formyl hydroxylamine compounds which are useful for inhibiting bacteria.

More specifically, the present invention is directed to a process for preparing a compound of the formula (VII)




comprising Step 1A:

Contacting a compound of the formula (I)




with a base in a suitable solvent to form the free base of compound (I), i.e., compound (II) of the formula (II)




followed by Step 1B:

Contacting compound (II) with a strong nucleophile/weak base in a suitable solvent under conditions to form compound (III) of the formula (III)




followed by Step 2A:

Contacting compound (III) with a formylating agent in a suitable solvent under conditions suitable to form a compound of formula (IV)




followed by Step 2B:

Contacting compound (IV) with an amine or an alkaline metal hydroxide in a suitable solvent under conditions to form a compound of formula (V)



followed by Step 3:

Contacting compound (V) with a compound of formula (VI)



in the presence of a suitable base and one or more coupling agents in a suitable solvent under conditions to form a compound of formula (VII)

wherein

Y is a hydroxy protecting group;

Each of R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub> and R<sub>5</sub> is, independently, hydrogen or alkyl, or (R<sub>2</sub> and R<sub>3</sub>) and/or (R<sub>4</sub> and R<sub>5</sub>) collectively form a C<sub>4-7</sub>cycloalkyl;

G is  $-O^{\ominus}\text{metal}^{\oplus}$  or  $-OH\cdot\text{amine}$ ;

X is  $-\text{CH}_2-$ ,  $-\text{S}-$ ,  $-\text{CH}(\text{OH})-$ ,  $-\text{CH}(\text{OR})-$ ,  $-\text{CH}(\text{SH})-$ ,  $-\text{CH}(\text{SR})-$ ,  $-\text{CF}_2-$ ,  $-\text{C}=\text{N}(\text{OR})-$  or  $-\text{CH}(\text{F})-$ ;  
wherein

R is alkyl;

$R_1$  is aryl or heteroaryl;


Z is a strong organic or inorganic acid; and

n is 0-3, provided that when n is 0, X is  $-\text{CH}_2-$ .

When the desired product is an N-oxide of an aromatic moiety having a nitrogen heteroatom (e.g., when  $R_1$  is Formula X, Xa or Xb), typically a pyridine derivative, it is necessary to perform an additional step after step 3, i.e., to oxidize the N of the aromatic ring (Step 4). Therefore, the present invention includes Step 4 which comprises contacting the compound of formula VII, wherein  $R_1$  is heteroaryl having an N heteroatom, with an oxidizing agent to form the corresponding N-oxide derivative.

In addition to the above process comprising Steps 1A-4 the present invention is directed to each of the steps individually, and to any two or more sequential steps.


In particular, the present invention provides a process for preparing intermediates useful in the preparation of a *N*-[1-oxo-2-alkyl-3-(*N*-hydroxyformamido)-propyl]- (carbonylamino-aryl or -heteroaryl)-azacyclo<sub>4,7</sub>alkane or thiazacyclo<sub>4,7</sub>alkane, e.g., a compound of formula (VIII)



wherein  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ , X and n are as defined above.


To convert the compound of formula (VII) to the compound of formula (VIII), the hydroxy protecting group is removed using conventional hydrogenolysis techniques known in the art, e.g., by contacting the compound of formula (VII) with a palladium catalyst, such as  $\text{Pd}/\text{BaSO}_4$  (see WO 02/102790 A1).

The  $R_1$  moiety can be a heteroaryl, e.g., an azacyclo<sub>4-7</sub>alkane, a thiazacyclo<sub>4-7</sub>alkane or an imidazacyclo<sub>4-7</sub>alkane. Specific examples of  $R_1$  moieties in the compounds disclosed herein are heteroaryls of formula (X)



wherein each of  $R_6$ ,  $R_7$ ,  $R_8$  and  $R_9$ , independently, is hydrogen, alkyl, substituted alkyl, hydroxy, alkoxy, acyl, acyloxy, SCN, halogen, cyano, nitro, thioalkoxy, phenyl, heteroalkylaryl, alkylsulfonyl or formyl.

A example of an  $R_1$  moiety is a heteroaryl of formula (Xa)



wherein  $R_6$ ,  $R_7$ ,  $R_8$  and  $R_9$  are as defined above for formula (X), e.g.,

wherein

- a)  $R_6$  is nitro, alkyl, substituted alkyl, phenyl, hydroxy, formyl, heteroalkylaryl, alkoxy, acyl or acyloxy; preferably alkyl, especially  $C_{1-7}$ alkyl; hydroxyl; or alkoxy, especially a  $C_{1-7}$ alkoxy; and  
 $R_7$ ,  $R_8$  and  $R_9$  are hydrogen; or
- b)  $R_6$ ,  $R_8$  and  $R_9$  are hydrogen; and  
 $R_7$  is alkyl, substituted alkyl, phenyl, halogen, alkoxy or cyano, preferably alkyl, especially  $C_{1-7}$ alkyl; substituted alkyl, especially substituted  $C_{1-7}$ alkyl, such as  $-CF_3$ ; or alkoxy, especially  $C_{1-7}$ alkoxy; or

- c)  $R_6$ ,  $R_7$  and  $R_9$  are hydrogen; and  
 $R_8$  is alkyl, substituted alkyl, halogen, nitro, cyano, thioalkoxy, acyloxy, phenyl, alkylsulfonyl or carboxyalkyl, preferably alkyl, especially  $C_{1-7}$ alkyl; substituted alkyl, especially  $-CF_3$ ; halogen; or carboxyalkyl; or
- d)  $R_6$ ,  $R_7$  and  $R_8$  are hydrogen; and  
 $R_9$  is alkyl, halogen or hydroxy; or
- e)  $R_7$  and  $R_9$  are hydrogen; and  
each of  $R_6$  and  $R_8$ , independently, is halogen, alkyl, substituted alkyl, phenyl or cyano; or
- f) Each of  $R_7$  and  $R_9$  is alkyl or substituted alkyl; and  
 $R_6$  and  $R_8$  are hydrogen; or
- g)  $R_6$  and  $R_9$  are hydrogen;  
 $R_7$  is alkyl or substituted alkyl; and  
 $R_8$  is nitro; or
- h)  $R_8$  and  $R_9$  are hydrogen;  
 $R_6$  is cyano; and  
 $R_7$  is alkoxy; or
- i)  $R_7$  and  $R_8$  are hydrogen; and  
 $R_6$  is alkyl, substituted alkyl, alkoxy or SCN; and  
 $R_9$  is alkyl or substituted alkyl; or
- j)  $R_6$  and  $R_7$  are hydrogen;  
 $R_8$  is nitro or halogen; and  
 $R_9$  is alkyl or substituted alkyl; or
- k)  $R_6$ ,  $R_7$ ,  $R_8$  and  $R_9$  are hydrogen; or
- l)  $R_6$  and  $R_7$  together with the carbon atoms to which they are attached form a phenyl group, preferably substituted with hydroxy; and  
 $R_8$  and  $R_9$  are hydrogen; or
- m)  $R_6$  and  $R_7$  are hydrogen; and  
 $R_8$  and  $R_9$  together with the carbon atoms to which they are attached form a phenyl group; or

n) n is 0; or

o) n is 0;

each of  $R_6$ ,  $R_7$ ,  $R_8$  and  $R_9$ , independently, is hydrogen, alkyl or halogen; and more particularly,  $R_6$ ,  $R_7$ ,  $R_8$  and  $R_9$  are hydrogen; or


p) n is 0;

$R_6$ ,  $R_8$  and  $R_9$  are hydrogen; and  $R_7$  is alkyl; or

q) n is 0;

$R_6$ ,  $R_7$  and  $R_9$  are hydrogen; and  $R_8$  is alkyl or halogen.

In another embodiment,  $R_1$  is of formula (Xb)




wherein

$R_6$ ,  $R_7$ ,  $R_8$  and  $R_9$  are as defined above for formula (X); in particular,  $R_7$  and  $R_8$  together with the carbon atoms to which they are attached form a phenyl group; and

$R_6$  and  $R_9$  are hydrogen.

In yet another embodiment, the  $R_1$  is of formula (XI)




wherein each of  $R_6$ ,  $R_7$ ,  $R_8$  and  $R_9$ , independently, is hydrogen, alkyl, substituted alkyl, phenyl, halogen, hydroxy or alkoxy, e.g.,

wherein



- a)  $R_6$  and  $R_8$  are hydrogen;  
 $R_9$  is hydrogen or alkyl; and  
 $R_7$  is alkyl, substituted alkyl or phenyl; or
- b)  $R_6$ ,  $R_7$  and  $R_9$  are hydrogen; and  
 $R_8$  is halogen, alkyl or substituted alkyl; or
- c)  $R_7$ ,  $R_8$  and  $R_9$  are hydrogen; and  
 $R_6$  is hydroxy.


In a particularly useful embodiment the heteroaryl is of the formula (XIa)



wherein  $R_6$ ,  $R_7$ ,  $R_8$  and  $R_9$  are as defined above for formula (XI).

In another embodiment,  $R_1$  is an unsubstituted phenyl or the phenyl is substituted with alkoxy, e.g., methoxy; or aryloxy, e.g., phenoxy.

In another embodiment, the  $R_1$  is of formula (XII)



wherein each of  $R_{10}$  and  $R_{11}$ , independently, is hydrogen or halogen. In particular,  $R_{10}$  and  $R_{11}$  are both either hydrogen or both halogen.

Unless otherwise stated, the following terms as used in the specification have the following meaning.

The term "cycloalkane" or "cycloalkyl" contains from 3- to 7-ring carbon atoms, and is, e.g., cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

The term "azacyclo<sub>4-7</sub>alkane" contains 1-ring heteroatom which is a nitrogen. It contains from 4-7, and especially 4- or 5-ring atoms including the heteroatom.

The term "thiazacyclo<sub>4-7</sub>alkane" contains 2-ring heteroatoms, nitrogen and sulfur. It contains from 4-7, and especially 5-ring atoms including the heteroatoms.

The term "imidazacyclo<sub>4-7</sub>alkane" contains 2-ring heteroatoms which are both nitrogen. It contains from 4-7, and especially 5-ring atoms including the heteroatoms.

The term "alkyl" refers to saturated or unsaturated aliphatic groups, such as alkenyl or alkynyl, cycloalkyl or substituted alkyl including straight-chain, branched-chain and cyclic groups having from 1-10 carbons atoms. Preferably "alkyl" or "alk", whenever it occurs, is a saturated aliphatic group or cycloalkyl, more preferably C<sub>1-7</sub>alkyl, particularly C<sub>1-4</sub>alkyl. Examples of "alkyl" or "alk" include, but are not limited to, methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, isobutyl, sec-butyl, *t*-butyl, *n*-pentyl, neopentyl, *n*-hexyl or *n*-heptyl, cyclopropyl and especially *n*-butyl.

The term "substituted alkyl" refers to an alkyl group that is substituted with one or more substituents preferably 1-3 substituents including, but not limited to, substituents, such as halogen, lower alkoxy, hydroxy, mercapto, carboxy, cycloalkyl, aryl, heteroaryl and the like. Examples of substituted alkyl groups include, but are not limited to, -CF<sub>3</sub>, -CF<sub>2</sub>-CF<sub>3</sub>, hydroxymethyl, 1- or 2-hydroxyethyl, methoxymethyl, 1- or 2-ethoxyethyl, carboxymethyl, 1- or 2-carboxyethyl and the like.

The term "aryl" or "Ar" refers to an aromatic carbocyclic group of 6-14 carbon atoms having a single ring including, but not limited to, groups, such as phenyl; or multiple condensed rings including, but not limited to, groups, such as naphthyl or anthryl; and is especially phenyl.

The term "heteroaryl" or "HetAr" refers to a 4- to 7-membered, monocyclic aromatic heterocycle or a bicycle that is composed of a 4- to 7-membered, monocyclic aromatic heterocycle and a fused-on benzene ring. The heteroaryl has at least one hetero atom, preferably one or two heteroatoms including, but not limited to, heteroatoms, such as N, O and S, within the ring. A preferred heteroaryl group is pyridinyl, pyrimidinyl or benzodioxolanyl.

The aryl or heteroaryl may be unsubstituted or substituted by one or more substituents including, but not limited to, C<sub>1-7</sub>alkyl, particularly C<sub>1-4</sub>alkyl, such as methyl, hydroxy, alkoxy, acyl, acyloxy, SCN, halogen, cyano, nitro, thioalkoxy, phenyl, heteroalkylaryl, alkylsulfonyl and formyl.

The term "carbonylamine", as used herein, refers to a -NHC(O)- group wherein the amino portion of the group is linked to the aryl/heteroaryl and the carbonyl portion of the group is linked to the azacyclo<sub>4-7</sub>alkane, thiaza<sub>4-7</sub>cycloalkane or imidazacyclo<sub>4-7</sub>alkane.

The term "heteroalkyl" refers to saturated or unsaturated C<sub>1-10</sub>alkyl as defined above, and especially C<sub>1-4</sub>heteroalkyl which contain one or more heteroatoms, as part of the main, branched or cyclic chains in the group. Heteroatoms may independently be selected from the group consisting of -NR-, where R is hydrogen or alkyl, -S-, -O- and -P-; preferably -NR-, where R is hydrogen or alkyl; and/or -O-. Heteroalkyl groups may be attached to the remainder of the molecule either at a heteroatom (if a valence is available) or at a carbon atom. Examples of heteroalkyl groups include, but are not limited to, groups, such as -O-CH<sub>3</sub>, -CH<sub>2</sub>-O-CH<sub>3</sub>, -CH<sub>2</sub>-CH<sub>2</sub>-O-CH<sub>3</sub>, -S-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>3</sub>, -CH<sub>2</sub>-CH(CH<sub>3</sub>)-S-CH<sub>3</sub> and -CH<sub>2</sub>-CH<sub>2</sub>-NH-CH<sub>2</sub>-CH<sub>2</sub>-.

The heteroalkyl group may be unsubstituted or substituted with one or more substituents, preferably 1-3 substituents including, but not limited to, alkyl, halogen, alkoxy, hydroxyl, mercapto, carboxy and, especially, phenyl. The heteroatom(s) as well as the carbon atoms of the group may be substituted. The heteroatom(s) may also be in oxidized form.

The term "alkoxy", as used herein, refers to a C<sub>1-10</sub>alkyl linked to an oxygen atom, or preferably C<sub>1-7</sub>alkoxy, more preferably C<sub>1-4</sub>alkoxy. Examples of alkoxy groups include, but are not limited to, groups, such as methoxy, ethoxy, *n*-butoxy, *tert*-butoxy and allyloxy.

The term "acyl", as used herein, refers to the group -(O)CR, where R is alkyl, especially C<sub>1-7</sub>alkyl, such as methyl. Examples of acyl groups include, but are not limited to, acetyl, propanoyl and butanoyl.

The term "acyloxy", as used herein, refers to the group -OC(O)R, wherein R is hydrogen, alkyl, especially C<sub>1-7</sub>alkyl, such as methyl or ethyl, or phenyl or substituted alkyl as defined above.

The term "alkoxycarbonyl", as used herein, refers to the group -COOR, wherein R is alkyl, especially, C<sub>1-7</sub>alkyl, such as methyl or ethyl.

The term "halogen" or "halo", as used herein, refers to chlorine, bromine, fluorine, iodine and, is especially, fluorine.


The term "thioalkoxy", as used herein, means a group -SR, where R is an alkyl as defined above, e.g., methylthio, ethylthio, propylthio, butylthio and the like.

The term "heteroalkylaryl", as used herein, means a heteroalkyl group, e.g., -O-CH<sub>2</sub>- substituted with an aryl group, especially, phenyl. The phenyl group itself may also be substituted with one or more substituents, such as halogen, especially, fluoro and chloro; and alkoxy, such as methoxy.

The term "alkylsulfonyl", as used herein, means a group -SO<sub>2</sub>R, wherein R is alkyl, especially, C<sub>1-7</sub>alkyl, such as methyl sulfonyl.


"Protecting group" refers to a chemical group that exhibits the following characteristics: 1) reacts selectively with the desired functionality in good yield to give a protected substrate that is stable to the projected reactions for which protection is desired; 2) is selectively removable from the protected substrate to yield the desired functionality; and 3) is removable in good yield by reagents compatible with the other functional group(s) present or generated in such projected reactions. Examples of suitable protecting groups may be found in Greene et al., *Protective Groups in Organic Synthesis*, 3<sup>rd</sup> Edition, John Wiley & Sons, Inc., NY (1999). Preferred hydroxy protecting groups include benzyl, Fmoc, TBDMS, photolabile protecting groups, such as Nvm, Mom and Mem. Other preferred protecting groups include NPEOC and NPEOM.

It will be appreciated that the compounds disclosed herein may exist in the form of optical isomers, racemates or diastereoisomers. In particular, in the compounds disclosed herein where R<sub>4</sub> and R<sub>5</sub> are different, the carbon atom to which the R<sub>4</sub> and R<sub>5</sub> groups are bonded is a chiral center and such compounds can exist in the *R*, *S* or racemic forms. It is contemplated that the process of the invention prepares the *R* optically pure form. By "optically pure" is meant that the enantiomeric purity is greater than 50%, preferably greater than 80%, more preferably greater than 90%, and most preferably greater than 95%. The optically pure *R* isomer of compound (I) can be used, in which case all subsequent compounds in the synthesis will remain in the *R* optically pure form, with respect to the same chiral carbon atom. Such *R* form of compound (I) is represented by former Ia below:




wherein  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  are as defined above. It is exemplified that in the compound of formula (I) that  $R_5$  is hydrogen and that  $R_4$  is  $C_{2-10}$ alkyl, more preferably  $C_{2-7}$ alkyl, and most preferably  $C_4$ alkyl.


It is further exemplified that in the optically pure compound of formula (I) that  $R_2$ ,  $R_3$ , and  $R_5$  are hydrogen and that  $R_4$  is alkyl; such a compound has the structure (Ib)




As an example, in compound (I),  $R_4$  is *n*-butyl, where such compound has the structure (Ic)



Further exemplified is that  $R_2$ ,  $R_3$  and  $R_5$  are hydrogen and that  $R_4$  is *n*-butyl; such compound has the structure (Id)



Alternatively, the racemate form of compound (I) can be used and then the *R* form can be resolved at a later step and the *R* form used for subsequent steps. For example, the compound formed after Step 3 or 3A, can be resolved into its *RS* and *SS* diastereomers and only the *RS* diastereomer used for subsequent steps. The *RS* diastereomer of compound (VII) is depicted below or formula (VIIa):



wherein  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $Y$ ,  $X$ ,  $R_1$  and  $n$  are as defined above, provided that  $R_4$  and  $R_5$  are different.

The optical isomers are resolved using standard techniques known in the art, for example, using silica gel column chromatography and an ethyl acetate/hexane solvent system. See, e.g., the methods taught in Chapter 4 of *Advanced Organic Chemistry*, 4<sup>th</sup> Edition, March, John Wiley and Sons, NY (1992).

In the compounds disclosed herein, the following significances are exemplified individually or in any sub-combination:

1.  $R_1$  is a heteroaryl of formula (Xa),

wherein

$R_6$ ,  $R_7$  and  $R_9$  are hydrogen and  $R_8$  is methyl or trifluoromethyl; or

$R_6$ ,  $R_7$  and  $R_9$  are hydrogen and  $R_8$  is fluoro; or

$R_6$ ,  $R_7$  and  $R_8$  are hydrogen and  $R_9$  is fluoro; or

$R_6$ ,  $R_8$  and  $R_9$  are hydrogen and  $R_7$  is ethyl or methoxy; or

$R_7$ ,  $R_8$  and  $R_9$  are hydrogen and  $R_6$  is hydroxy; or

$R_7$  and  $R_8$  are hydrogen,  $R_6$  is methoxy and  $R_9$  is methyl; or

$R_1$  is a heteroaryl of formula (Xb),

wherein

$R_6$ ,  $R_7$  and  $R_9$  are hydrogen and  $R_8$  is fluoro or trifluoromethyl; or

$R_6$ ,  $R_8$  and  $R_9$  are hydrogen and  $R_7$  is ethyl; preferably  $R_1$  is a heteroaryl of formula (Xa),

wherein  $R_6$ ,  $R_8$  and  $R_9$  are hydrogen and  $R_7$  is ethyl or a heteroaryl of formula (Xb),

wherein  $R_6$ ,  $R_7$  and  $R_9$  are hydrogen and  $R_8$  is fluoro.

2. X is  $-\text{CH}_2-$ ,  $-\text{CH}(\text{OH})-$ ,  $-\text{CH}(\text{OR})-$ ,  $-\text{CF}_2-$  or  $-\text{CH}(\text{F})-$ , preferably X is  $-\text{CH}_2-$ ;
3.  $R_4$  is alkyl, preferably  $\text{C}_{1-7}$  alkyl, such as *n*-butyl;
4. *n* is 1.

Temperature and pressure are not known to be critical for carrying out any of the steps of the invention, i.e., Steps 1A-4. Generally, for any of the steps, a temperature of about  $-10^\circ\text{C}$  to about  $150^\circ\text{C}$ , typically about  $0^\circ\text{C}$  to about  $80^\circ\text{C}$ , is employed. Typically about atmospheric pressure is used for convenience; however, variations to atmospheric pressure are not known to be detrimental. Oxygen is not known to be detrimental to the process, therefore for convenience the various steps can be performed under ambient air, although an inert atmosphere, such as nitrogen or argon, can be used if desired. For convenience equimolar amounts of reactants or reagents, as appropriate, are typically used; however molar ratios can vary from about 1 to 2 equivalents, relative to the other reactant/reagent. The pH for the various steps is typically about 2 to about 12. The solvent used for the various steps are typically organic solvents, although in some situations aqueous/organic solvents can be used. Examples of suitable solvents include dioxane; methylene chloride; dichloromethane; toluene, acetone; methyl ethyl ketone; THF; isopropyl acetate; DMF; alcohols, especially, ethyl acetate, acetonitrile, higher-branched alcohols, such as *t*-butanol; and the like.

For Step 1A, a typical temperature is about 10°C to about 40°C, more typically about 15°C to about 25°C; and a typical reaction time is about 0.1 hours to about 3 hours, more typically about 0.25 hours to about 1 hour. A pH of about 6 pH to about 10 pH, typically about 8 pH to about 9 pH, more typically about 9 pH, is employed. The base for Step 1A is a water soluble base such as sodium carbonate, sodium bicarbonate, potassium carbonate, potassium bicarbonate, an alkaline metal hydroxide, e.g., sodium hydroxide, potassium hydroxide, and the like. The solvent for Step 1A is a biphasic solvent, i.e., a mixture of water and an organic solvent immiscible with water, for example, ethyl acetate, methylene chloride, diethyl ether, methyl *t*-butyl ether, isopropyl acetate, and the like. An example of a solvent is water/ethyl acetate. To prepare the starting compound of formula (I) for Step 1A (i.e., a salt) a strong acid is added to the corresponding free amine in solution with an organic solvent such as ethyl acetate, ethyl ether, and the like. The Z substituent, i.e., the strong acid, must be of sufficient strength to form a salt of the amine which results in the compound of formula (I) precipitating from the organic solution. The Z substituent is a strong organic or inorganic acid such as HCl, HBr, benzenesulfonic acid, toluenesulfonic acid, camphorsulfonic acid, and the like.


For Step 1B, a typical temperature is about -10°C to about 10°C, more typically about -3°C to about 2°C; and a typical reaction time is about 0.5 hours to about 5 hours more typically about 0.75 hours to about 1.5 hours. The pH for Step B is typically about 8 pH to about 11 pH. The strong nucleophile/weak base used in Step 1B can be, for example, lithium hydroperoxide or a thiolate salt of an alkaline metal such as the sodium salt of propanethiol. The strong nucleophile/weak base is typically formed *in situ*, such as by adding hydrogen peroxide and an alkaline metal hydroxide, for example adding hydrogen peroxide and lithium peroxide to form lithium hydroperoxide *in situ*. The solvent for Step 2A can be a mixture of water and an ether solvent that is water miscible, such as THF, dimethylethane, dioxane, and the like. A typical solvent is THF/water.

For Step 2A, a typical temperature is about -20°C to about 20°C, more typically about -10°C to about 5 °C; and a typical reaction time is about 0.25 hours to about 2 hours, more typically about 0.3 hours to about 1 hour. The pH for Step 2A is typically, about 1 pH to about 6 pH. The formylating agent for Step 2A is typically formed *in situ*, such as by adding formic acid and acetic anhydride to form formic acetic anhydride. The solvent for Step 2A is an inert solvent in which the desired compound is soluble, for example, ethyl acetate,



isopropyl acetate, methyl acetate, n-butyl acetate and the like. A typical solvent is ethyl acetate.

For Step 2B, a typical temperature is about -5°C to about 40°C, more typically about 15°C to about 25°C; and a typical reaction time is about 1 hour to about 5 hours, more typically about 2 hours to about 3 hours. The pH for Step 2B is typically about 1 pH to about 6 pH. Typical solvents for Step 2B include ethyl acetate, iso-propyl acetate, heptane, and the like. A particular example of a solvent is heptane. Examples of G substituents include -O<sup>o</sup>metal<sup>o</sup> wherein the metal is Na, K, Mg, Li, or -OH•amine wherein the amine of the formula HNR'R', wherein each R' is a straight chain, branched chain or cyclo alkyl group of 1 to 8 carbon atoms, more typically 1 to 6 carbon atoms. A typical example of a G substituent is -OH•amine wherein the amine is dicyclohexylamine. Therefore, an example of the compound of formula (V) has the structure:



For Step 3, a typical temperature is about 10°C to about 40°C, more typically about 15°C to about 25°C; and a typical reaction time is about 5 minutes to about 15 hours, more typically about 10 minutes to about 10 hours. The pH for Step 3 is typically about 5 to about 9. The solvent for Step 3 is a biphasic solvent, i.e., a mixture of water and an organic solvent immiscible with water, for example, ethyl acetate, methylene chloride, diethyl ether, methyl t-butyl ether, isopropyl acetate, and the like. A typical solvent is water/ethyl acetate. Typical bases for Step 3 include tertiary amine bases such as N-methylmorpholine, triethylamine, diisopropylethylamine, and the like. The coupling agent can be a conventional coupling agent known in the art, for example as disclosed in J. Jones, "The Chemical Synthesis of Peptides", Clarendon, Oxford, 1991 and P. Lloyd Williams, F. Albericio and E. Girault, Tetrahedron, 1993, 49, 11065, incorporated herein by reference. One or more coupling agents are used. Examples of coupling agents include EDCI, HOBt, DCC, HATU,

BOP, FDPP, cross linked enzyme crystals such as PEPTI CLEC-TR, and the like. A typical coupling agent is EDCI/HOBt. A typical molar ration of DCCI:HOBt is about 1:5 to about 5:1.

For Step 4, a typical temperature is about 10°C to about 35°C, more typically about 20°C to about 22°C; and a typical reaction time is about 60 minutes to about 18 hours, more typically about 4 hours to about 8 hours. The pH for Step 4 is typically about 4 to about 8. The solvent for Step 4 is typically an organic solvent, i.e., ethyl acetate, iso-propyl acetate, methylene chloride, and the like. The oxidizing agent can be a conventional agent known in the art, for example as disclosed in March, "Advanced Organic Chemistry", 5th Ed., Wiley Interscience, NY, Chapter 19, incorporated herein by reference. Typical oxidizing agents include urea/hydrogen peroxide with phthalic anhydride; magnesium monoperoxyphthalate; MCPBA, Oxone (available from Aldrich), and the like.

Insofar as the production of starting materials is not particularly described, the compounds are known or may be prepared analogously to methods known in the art or as disclosed in the examples hereinafter.

The following abbreviations are used:

Ac = acetyl

BOP = [benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate

CDMT = chlorodimethoxy triazine

DIEA = diisopropylethylamine

DCC = dicyclohexylcarbodiimide

DMF = dimethylformamide

EDCI = 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride

2-EHA = 2-ethylhexanoic acid

EtOAc = ethyl acetate

EtOH = ethanol

HATU = [O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate]  
isobutyl chloroformate

HPLC = high performance liquid chromatography

MCPBA = metachloroperoxybenzoic acid

MeOH = methanol

MMPP = magnesium monoperoxyphthalate


RT = room temperature

THF = tetrahydrofuran

The following illustrates a preferred process of the invention.

**Reaction Scheme**

The following examples illustrate the invention but should not be interpreted as a limitation thereon. Product numbers refer to the preferred reaction scheme depicted immediately below.



General procedure for the synthesis of intermediates useful for the preparation of:

**(2S)-N-(5-fluoro-1-oxido-2-pyridinyl-1-[(2R)-2-[formylhydroxyamino)methyl]-1-oxohexyl]-2-pyrrolidinecarboxamide, magnesium salt**

**Step 1: (2R)-2-[[[(phenylmethoxy)amino]methyl]-hexanoic acid (A8)**

To a solution of the *p*-TSA salt (**A7**) (58.3 g, 0.1 mol) in ethyl acetate (200 mL) and water (50 mL) was added 1 N Na<sub>2</sub>CO<sub>3</sub> (185 mL). The two phase mixture was stirred for 15 minutes at RT and the lower aqueous layer was separated. The organic layer was washed with water (2 x 50 mL), and concentrated to give the free base of **A7**.

The **A7** free base (41.0 g, 0.1 mol) was dissolved in THF (395 mL) and water (107 mL) and cooled to -3°C. To this solution was added 30% hydrogen peroxide (26.1 g, 0.23 mol) keeping the temperature at -3°C. In a separate flask, a solution of lithium hydroxide (5.0 g, 0.12 mol) in water (107 mL) was prepared and added slowly to the **A7**/hydrogen peroxide solution keeping the temperature at -3°C. The mixture was stirred for 45 minutes at this temperature.

A solution of sodium sulfite (43.5 g, 0.345 mol) in water (855 mL) was added slowly keeping the temperature below 10°C and the reaction mixture was allowed to warm to RT. The solution was partially concentrated under vacuum to remove the THF and the aqueous portion was extracted with ethyl acetate (6 x 110 mL). The aqueous portion was then acidified with 3 N HCl (78 mL) and extracted with ethyl acetate (2 x 215 mL). The ethyl acetate extracts were combined and washed with water (2 x 110 mL). The organic solution was partially concentrated under vacuum (200 mL) to give a colorless solution of **A8** which was used as is in the following step.

A sample was concentrated completely for characterization.

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.4 (s, 5H), 6.85 (bs, 2H), 4.75 (dd, 2H), 3.1 (m, 2H), 2.8 (m, 1H), 2.7 (m, 1H), 2.55 (m, 1H), 1.2 (m, 4H), 0.88 (m, 3H). ES-MS: calcd. for C<sub>14</sub>H<sub>21</sub>NO<sub>3</sub> (251.3); found: 252.2 [M+H].

Step 2: (2*R*)-2-[[formyl(phenylmethoxy)amino]methyl]-hexanoic acid dicyclohexylamine salt (**A10**)

Acetic anhydride (15.3 g, 0.15 mol) was cooled to 0-5°C and treated with 96% formic acid (27.6 g, 0.6 mol) keeping the temperature below 10°C. The mixture was stirred for 15 minutes at 0-5°C and then warmed to RT and stirred for 15 minutes more.

In a second flask, **A8** ethyl acetate solution (502 g, 0.75 mol) was cooled to -15°C and the formic acid/acetic anhydride mixture was added to it keeping the temperature at -10 ± 5°C. The reaction mixture was stirred for 20 minutes at this temperature and then water (5.4 g) was added. After stirring for 15 minutes, the solution was warmed to RT. The solution was

concentrated under vacuum (final volume = 70-90 mL). Toluene (240 mL) was added and the solution was again concentrated under vacuum (final volume = 70-80 mL).

In a separate flask a mixture of dicyclohexylamine (16.3 g) in heptane (240 mL) was prepared and this was added to the concentrate at RT. The mixture was seeded and held with stirring for 2 hours. Heptane (145 mL) was added and the suspension was held for 8 hours at RT. The solids were isolated by filtration and dried under vacuum to give the title compound.

m.p.: 83-86°C; <sup>1</sup>H NMR (CDCl<sub>3</sub>, rotamers): δ 8.05 (bd, 1H), 7.3-7.65 (m, 5H), 4.75-5.1 (m, 2H), 3.5-4.0 (m, 2), 3.1-3.39 (m, 1H), 2.9 (m, 3H), 2.65 (m, 1H), 1.0-2.15 (m, 26H), 0.9 (s, 3H). ES-MS: calcd. for C<sub>15</sub>H<sub>21</sub>NO<sub>4</sub> (free acid) (279); found: 280.1 [M+H].

Step 3: (2S)-N-(5-fluoro-2-pyridinyl)-1-[(2R)-2-[[formyl(phenylmethoxy)amino]methyl]-1-oxohexyl]-2-pyrrolidinecarboxamide (A11)

A solution of A10 (34.55 g, 75 mmol) in ethyl acetate (300 mL) was mixed with a citric acid solution (30 g of citric acid in 270 mL of water) and stirred at RT for 10 minutes. The layers were separated and the upper organic layer was washed with water (2 x 225 mL). At this point, N-(5-fluoro-2-pyridinyl)-(2S)-2-pyrrolidinecarboxamide dihydrobromide (33.39 g, 90 mmol) was added followed by water (60 mL) and HOBt (12.81 g, 82.5 mmol).

The mixture was cooled to 0-5°C and EDCI (40.26 g, 210 mmol) and water (60 mL) were added. This was followed by the addition of N-methylmorpholine (47.79 g, 472.5 mmol). The reaction was stirred at RT overnight.

The lower aqueous layer was separated and the upper organic layer was washed with water (4 x 225 mL). The organic layer was filtered through a column of silica gel (83.4 g) and the column was further eluted with an additional volume of ethyl acetate (3 x 41 mL). The suitable fractions were combined and concentrated under vacuum to a specific volume (225 mL).

This solution was warmed to 50°C and treated with heptane (675 mL). The solution was then cooled to 45°C and seeded. The slurry was cooled to below -10°C and held for 2 hours. The solids were isolated by filtration and dried under vacuum to give the title compound.

m.p.: 98°C;  $^1\text{H}$  NMR (DMSO, rotamers):  $\delta$  10.6, 10.8 (s, 1H), 8.2 (s, 1H), 7.5-8.2 (m, 3H), 6.95-7.4 (m, 5H), 4.8 (s, 2H), 4.55 (bs, 1H), 3.2-3.8 (m, 4H), 2.9 (bs, 1H), 1.6-2.4 (m, 4H), 1.0-1.55 (m, 6H), 0.8 (s, 3H). ES-MS: calcd. for  $\text{C}_{25}\text{H}_{31}\text{FN}_4\text{O}_4$  (470.6); found: 471.2 [M+H], 493.2 [M+Na].

Step 4: (2S)-N-(5-fluoro-1-oxido-2-pyridinyl)-1-[(2R)-2-  
[[formyl(phenylmethoxy)amino]methyl]-1-oxohexyl]-2-pyrrolidinecarboxamide (A12)

A mixture of magnesium monoperoxyphthalate (69.25 g, 140 mmol) in water (128 mL) and isopropyl acetate (300 mL) was stirred and a solution of A11 (32.94 g, 70 mmol) in isopropyl acetate (162 mL) was added. The mixture was stirred for 17 hours at RT.


The bottom aqueous layer was separated and a solution of sodium sulfite (8.82 g, 70 mmol) in water (160 mL) was added. After stirring for 20 minutes, the bottom aqueous layer was separated and sodium carbonate (20 g, 190 mmol) in water (300 mL) was added. After stirring for 20 minutes, the bottom aqueous layer was separated and a solution of sodium chloride (19.0 g) in water (131 mL) was added. The layers were separated and the organic layer was concentrated under vacuum to a final volume of 92 mL.

The solution was filtered and the filtrate was heated to 40°C and heptane (80 mL) was added. The solution was allowed to slowly cool to 30°C and seed crystals were added. The mixture was held for one hour at this temperature and then cooled to 22°C and more heptane was added (545 mL). After all of the heptane was added, the suspension was held at 22°C for 2 hours and then further cooled to below -10°C and held for 1 hour. The solids were isolated by filtration and dried under vacuum to give the title compound.

m.p.: 70°C;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , rotamers):  $\delta$  10.35 (s, 1H), 8.45-8.75 (m, 1H), 7.61-8.45 (m, 2H), 7.35 (s, 5H), 7.05 (s, 1H), 4.65-5.22 (m, 2H), 4.1-4.65 (m, 1H), 3.25-4.1 (m, 3.5H), 2.64-3.2 (m, 1.5H), 1.02-2.42 (m, 10H), 0.85 (s, 3H). ES-MS: calcd. for  $\text{C}_{25}\text{H}_{31}\text{FN}_4\text{O}_5$  (486.5); found: 487.2 [M+H].


## WHAT IS CLAIMED IS:

1. A process for preparing a compound of the formula (VII)




comprising Step 1A:

contacting a compound of the formula (I)




with a base in a suitable solvent to form the free base of compound (I), i.e., compound (II) of the formula (II)



followed by Step 1B:




contacting compound (II) with a strong nucleophile/weak base in a suitable solvent under conditions to form compound (III) of the formula (III)




followed by Step 2A:

contacting compound (III) with a formylating agent in a suitable solvent under conditions suitable to form a compound of formula (IV)




followed by Step 2B:

contacting compound (IV) with an amine or an alkaline metal hydroxide in a suitable solvent under conditions to form a compound of formula (V)



followed by Step 3:

contacting compound (V) with a compound of formula (VI)



in the presence of a suitable base and one or more coupling agents in a suitable solvent under conditions to form a compound of formula (VII)

wherein

Y is a hydroxy protecting group;

each of R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub> and R<sub>5</sub> is, independently, hydrogen or alkyl, or (R<sub>2</sub> and R<sub>3</sub>) and/or (R<sub>4</sub> and R<sub>5</sub>) collectively form a C<sub>4-7</sub> cycloalkyl;

G is  $-O^{\ominus}\text{metal}^{\oplus}$  or  $-OH\cdot\text{amine}$ ;

X is  $-\text{CH}_2-$ ,  $-\text{S}-$ ,  $-\text{CH}(\text{OH})-$ ,  $-\text{CH}(\text{OR})-$ ,  $-\text{CH}(\text{SH})-$ ,  $-\text{CH}(\text{SR})-$ ,  $-\text{CF}_2-$ ,  $-\text{C}=\text{N}(\text{OR})-$  or  $-\text{CH}(\text{F})-$ ;

R is alkyl;


$\text{R}_1$  is aryl or heteroaryl;

Z is a strong organic or inorganic acid; and

n is 0-3, provided that when n is 0, X is  $-\text{CH}_2-$ .

2. The process of Claim 1 followed by Step 4, contacting the compound of formula VII, wherein  $\text{R}_1$  is heteroaryl having an N heteroatom, with an oxidizing agent to form the corresponding N-oxide derivative.

3. The process of Claim 2 followed by the additional step of removing the hydroxyl protecting group of compound VII to form the compound of formula VIII:



wherein  $\text{R}_1$ ,  $\text{R}_2$ ,  $\text{R}_3$ ,  $\text{R}_4$ ,  $\text{R}_5$ , X and n are as defined above.

4. The process of Claim 1, wherein

each of  $\text{R}_2$ ,  $\text{R}_3$  and  $\text{R}_5$  is hydrogen;


$\text{R}_4$  is butyl;

X is  $-\text{CH}_2-$ ;

n is 1;

Y is benzyl or *t*-butyldimethylsilyl; and

$\text{R}_1$  is of the formula



wherein


$\text{R}_6$  and  $\text{R}_9$  are hydrogen;

$R_7$  is hydrogen or  $C_{1-7}$ alkyl; and  
 $R_8$  is hydrogen, halogen or  $C_{1-7}$ alkyl.

5. The process of Claim 4,  
 wherein

$R_7$  is hydrogen; and  
 $R_8$  is fluoro.

6. The process of claim 1, wherein  $R_1$  is of the formula (Xla)



each of  $R_2$ ,  $R_3$  and  $R_5$  is hydrogen;

$R_4$  is butyl;

$X$  is  $-CH_2-$ ;

$n$  is 1;

$Y$  is benzyl or t-butyldimethylsilyl;

$R_6$  and  $R_9$  are hydrogen;

$R_7$  is hydrogen or  $C_{1-7}$ alkyl; and

$R_8$  is hydrogen, halogen or  $C_{1-7}$ alkyl.

7. The process of Claim 6 wherein  $R_8$  is halo or ethyl.
8. The process of Claim 6 wherein  $R_7$  is hydrogen and  $R_8$  is fluoro.
9. The process of Claim 1 wherein

for Step 1A the temperature is about  $10^\circ\text{C}$  to about  $40^\circ\text{C}$ , the water soluble base is sodium carbonate, sodium bicarbonate, potassium carbonate, potassium bicarbonate, or an alkaline metal hydroxide, and the solvent is water/ethyl acetate,

for Step 1B the temperature is about  $-10^{\circ}\text{C}$  to about  $10^{\circ}\text{C}$ , the strong nucleophile/weak base is lithium hydroperoxide, and the solvent is THF/water,

for Step 2A the temperature is about  $-20^{\circ}\text{C}$  to about  $20^{\circ}\text{C}$ , the formylating agent is formic acetic anhydride, and the solvent is ethyl acetate,


for Step 2B the temperature is about  $-5^{\circ}\text{C}$  to about  $40^{\circ}\text{C}$ , the solvent is heptane and the G substituent is of the formula  $-\text{OH}\cdot\text{amine}$  wherein the amine is dicyclohexylamine,

for Step 3 the temperature is about  $10^{\circ}\text{C}$  to about  $40^{\circ}\text{C}$  the solvent is water/ethyl acetate, and the coupling agent is EDCI/HOBt, and


for Step 4 the temperature is about  $10^{\circ}\text{C}$  to about  $35^{\circ}\text{C}$ , the solvent is ethyl acetate and the oxidizing agent is urea/hydrogen peroxide with phthalic anhydride or magnesium monoperoxyphthalate.

10. A process comprising

contacting a compound of the formula (I)



with a base in a suitable solvent to form compound (II) of formula



wherein


Y is a hydroxy protecting group;

each of  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  is, independently, hydrogen or alkyl, or ( $R_2$  and  $R_3$ ) and/or ( $R_4$  and  $R_5$ ) collectively form a  $C_{4-7}$ cycloalkyl;


and Z is a strong organic or inorganic acid.

11. A process comprising

contacting compound (II) of the formula



with a strong nucleophile/weak base in a suitable solvent under conditions to form compound (III) of the formula




wherein

Y is a hydroxy protecting group; and


each of  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  is, independently, hydrogen or alkyl, or ( $R_2$  and  $R_3$ ) and/or ( $R_4$  and  $R_5$ ) collectively form a  $C_{4-7}$ cycloalkyl.

12. A process comprising

contacting compound (III) of the formula



with a formylating agent in a suitable solvent under conditions suitable to form a compound of formula (IV)




wherein

Y is a hydroxy protecting group; and


each of  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  is, independently, hydrogen or alkyl, or ( $R_2$  and  $R_3$ ) and/or ( $R_4$  and  $R_5$ ) collectively form a  $C_{4-7}$ cycloalkyl.

13. A process comprising

contacting compound (IV) of the formula



with an amine or an alkaline metal hydroxide in a suitable solvent under conditions to form a compound of formula (V)




wherein


Y is a hydroxy protecting group;

each of  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  is, independently, hydrogen or alkyl, or ( $R_2$  and  $R_3$ ) and/or ( $R_4$  and  $R_5$ ) collectively form a  $C_{4-7}$ cycloalkyl; and  
 $G$  is  $-O^{\ominus}\text{metal}^{\oplus}$  or  $-OH\cdot\text{amine}$ .


14. A process comprising  
 contacting compound (V) of the formula



with a compound of formula (VI)



in the presence of a suitable base and one or more coupling agents in a suitable solvent  
 under conditions to form a compound of formula (VII)



wherein

$Y$  is a hydroxy protecting group;

each of  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  is, independently, hydrogen or alkyl, or ( $R_2$  and  $R_3$ ) and/or ( $R_4$  and  $R_5$ ) collectively form a  $C_{4-7}$ cycloalkyl;

$G$  is  $-O^{\ominus}\text{metal}^{\oplus}$  or  $-OH\cdot\text{amine}$ ;

X is  $-\text{CH}_2-$ ,  $-\text{S}-$ ,  $-\text{CH}(\text{OH})-$ ,  $-\text{CH}(\text{OR})-$ ,  $-\text{CH}(\text{SH})-$ ,  $-\text{CH}(\text{SR})-$ ,  $-\text{CF}_2-$ ,  $-\text{C}=\text{N}(\text{OR})-$  or  $-\text{CH}(\text{F})-$ ;

R is alkyl;

$\text{R}_1$  is aryl or heteroaryl; and

n is 0-3, provided that when n is 0, X is  $-\text{CH}_2-$ .



# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP2004/006915

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D401/12

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 02/102790 A (NOVARTIS AG ; GARCIA ALVAREZ SALVADOR (US); JAIN RAKESH K (US); JACOBS) 27 December 2002 (2002-12-27) cited in the application page 14 - page 18 abstract	10
X	US 6 503 897 B1 (SPAVOLD ZOE MARIE ET AL) 7 January 2003 (2003-01-07) Scheme 1, Steps 4 and 5 abstract	11, 12
X	US 6 423 690 B1 (DAVIES STEPHEN JOHN ET AL) 23 July 2002 (2002-07-23) Scheme 1, Step C column 18, lines 8-21 abstract	12

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*G\* document member of the same patent family

Date of the actual completion of the international search

11 November 2004

Date of mailing of the international search report

24/11/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Guspanova, J

# INTERNATIONAL SEARCH REPORT

Form PCT/ISA/210 (patent family members)

International Application No

PCT/EP/006915

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
WO 02102790	A	27-12-2002	BR	02 10377 A	10-08-2004
			CA	24 48526 A1	27-12-2002
			CZ	200 33388 A3	17-03-2004
			WO	021 02790 A1	27-12-2002
			EP	14 01828 A1	31-03-2004
			HU	04 00208 A2	28-06-2004
			SK	152 42003 A3	07-07-2004
			US	20030 45479 A1	06-03-2003
			US	20032 10101 A1	13-11-2003
US 6503897	B1	07-01-2003	AU	35 65700 A	16-10-2000
			EP	11 65532 A1	02-01-2002
			WO	00 58294 A1	05-10-2000
			JP	20025 40197 T	26-11-2002
US 6423690	B1	23-07-2002	AT	2 66394 T	15-05-2004
			AU	7 49699 B2	04-07-2002
			AU	25 29299 A	23-08-1999
			BR	99 07689 A	14-11-2000
			CA	23 20476 A1	12-08-1999
			CN	12 98299 T	06-06-2001
			DE	699 17221 D1	17-06-2004
			DK	10 52984 T3	20-09-2004
			EP	10 52984 A1	22-11-2000
			GB	23 49884 A	15-11-2000
			WO	99 39704 A1	12-08-1999
			HU	01 02901 A2	28-12-2001
			JP	20025 02815 T	29-01-2002
			NO	200 03969 A	28-09-2000
			NZ	5 05675 A	26-11-2002
			NZ	5 21033 A	31-10-2003
			PL	3 42296 A1	04-06-2001
			PT	10 52984 T	30-09-2004
			TR	2000 02311 T2	21-11-2000
			US	20021 65167 A1	07-11-2002

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**